



# Principles of Interferometry and Science Results at the CHARA Array



Gail Schaefer

The CHARA Array of  
Georgia State University

Mount Wilson, CA




# Principles of Interferometry






# Interferometer



Resolution ~ 0.5 mas for 300 meter baseline in the H-band ( $1.6 \mu\text{m}$ )




# Fringe Visibility






- Amplitude of fringes = Visibility
  - Point Source:  $V = 1.0$
  - Resolved source: loss of coherence reduces fringe visibility
  - Measures the size and geometry of source

# Fringe Visibility



- The visibility is the Fourier Transform of the brightness distribution
- Analytic functions for simple geometries
- Berger & Segransan  
“Introduction to visibility modeling” 2007, New Ast Rev, 51, 576



# Angular Diameters



- Visibility amplitude
  - size and structure of source





# Elliptical Disk






# Elliptical Disk






# Surface Features






# Surface Features



# Binary Stars


## Separated Fringe Packet Binaries



Farrington et al. (2010)


# Binary Stars

## Visibility Modulation



- Fringe packets for the two components overlap
- Fringe visibility varies periodically
  - binary separation
- Minimum in curve
  - flux ratio =  $\frac{1 - V_{min}}{1 + V_{min}}$

# Closure Phase



Monnier, "Phases in Interferometry" 2007, New Ast Rev, 51, 604



- Atmosphere corrupts phase information at vis/IR wavelengths
- Closure phase (3 or more telescopes):
  - $CP = \phi_{12} + \phi_{23} + \phi_{31}$
- Cancels atmospheric effects
- Point symmetric object will have closure phase of  $0^\circ$  or  $180^\circ$
- Measures asymmetries in source distribution





# Binary Stars

Visibility (S1-E1)

Closure Phase (S1-E1-W1)




Flux ratio = 0.99



Flux ratio = 0.50

# Differential Visibilities and Phases



- Spectrally dispersed interferometry
  - emission lines (BrG, Ha)
  - velocity structure

- Drop in visibility across emission line
  - variation in size and flux ratio between star and disk
- “S” shaped profile in differential phase
  - photo-center shift across wavelength channels



# Interferometric Observables

- Visibility amplitude
  - size and structure of source
- Closure phase
  - asymmetries in source distribution
- Differential visibilities and phases
  - emission lines
  - velocity structure






# UV Coverage

$$u = B_x / \lambda$$

$$v = B_y / \lambda$$





- Interferometer baseline projected on to the plane of the sky
- Position angle and projected baseline length will change as the earth rotates

# UV Coverage

$$u = B_x / \lambda$$

$$v = B_y / \lambda$$



- Interferometer baseline projected on to the plane of the sky
- Position angle and projected baseline length will change as the earth rotates



# Science Review: Recent Results at the CHARA Array






# Outline

- Stellar Astrophysics
  - Stellar Diameters
  - Rapid Rotation
  - Spotted Stars
- Binary Stars
  - Orbits
  - High Contrast Binaries
  - Interacting Binaries
- Circumstellar Disks
  - Be Stars
  - Young Stellar Objects
- Transient Events
  - Nova Explosions



# Stellar Diameters




- Empirical HRD
- ~ 290 stars
- $\sigma_{\theta} < 5\%$
- Derive empirical color-temperature relations



# Fundamental Stellar Parameters



- Angular diameter + parallax
  - Linear radius





# Fundamental Stellar Parameters

- Angular diameter + parallax
  - Linear radius
- Spectral Energy Distribution
  - Bolometric flux







# Fundamental Stellar Parameters



- Angular diameter + parallax
  - Linear radius
- Spectral Energy Distribution
  - Bolometric flux
- Effective Temperature

- $$F_{bol} = \frac{1}{4} \theta^2 \sigma T^4$$







# Ages of Stars



- Diameters of 7 A-type stars in Ursa Major moving group
- Compare with evolutionary models that include rotation (mass, age)
- Age =  $414 \pm 23$  Myr
- Jones et al. 2015




# Ages of Stars



Ligi et al. 2016

- Diameters of 18 bright exoplanet host stars and candidates
- Compare (R, Teff) with evolutionary tracks to compute mass and age estimates
- Typically, two distinct solutions (old and young age)


# Exoplanet Host Stars



- Size of habitable zones
  - $L$ ,  $T_{\text{eff}}$
- Age and mass of host star
- Physical parameters of planets
  - Radius of transiting planets



# Asteroseismology



Mass, radius, mean density, and surface gravity (need  $T_{\text{eff}}$ )

$$\nu_{\text{max}} \propto (M / R^2) (T_{\text{eff}})^{-0.5}$$


$$\langle v_{nl} \rangle \propto \langle \rho \rangle^{0.5}$$

Oscillation power spectrum

$\langle v_{nl} \rangle$ : frequency separation of modes

$\nu_{\text{max}}$ : frequency of maximum power

# Asteroseismology




Mass, radius, mean density, and surface gravity (need  $T_{\text{eff}}$ )

$$\nu_{\text{max}} \propto (M / R^2) (T_{\text{eff}})^{-0.5}$$

$$\langle \nu_{\text{nl}} \rangle \propto \langle \rho \rangle^{0.5}$$



Oscillation power spectrum  
 $\langle \nu_{\text{nl}} \rangle$ : frequency separation of modes  
 $\nu_{\text{max}}$ : frequency of maximum power



Test asteroseismic scaling relations for main sequence stars

Huber et al. (2012)

# Cepheids





- Radial velocity and angular diameter variation of delta Cephei measured over the pulsational phase (Merand et al. 2005)
- Improve calibration of Baade-Wesselink technique for determining pulsation parallaxes




# Cepheids - Merand et al. 2015

delta Cep (P~5.4d) p=1.288 d=274.0pc E(B-V)=0.032  $K_{cep}=0.025\text{mag}$   $H_{cep}=0.020\text{mag}$



- Integrated parallax of pulsation method
- Simultaneous time series fit:
  - Photometry
  - Spectroscopy
  - Interferometry
- Mitigate systematics:
  - projection factor
- 2% accuracy on radius and distance


# Rapid Rotators




- Oblateness
- Gravity darkening
  - $T_{\text{eff}} \sim g^{\beta}$




# Rapid Rotators




- Oblateness
- Gravity darkening
  - $T_{\text{eff}} \sim g^{\beta}$
  - von Zeipel model:  $\beta = 0.25$
  - empirically derived  $\beta = 0.19$



# Spotted Stars




- Magnetically active star zeta Andromedae
- Rotation Period: 18 days
- $\theta = 2.502 \pm 0.008$  mas




Roettenbacher et al. (2016)

# Spotted Stars




- Magnetically active star zeta Andromedae
- Rotation Period: 18 days
- $\theta = 2.502 \pm 0.008$  mas
- Direct confirmation of persistent polar spot




Roettenbacher et al. (2016)

# Spotted Stars

2011



2013




- Magnetically active star zeta Andromedae
- Rotation Period: 18 days
- $\theta = 2.502 \pm 0.008$  mas
- Direct confirmation of persistent polar spot
- Transient lower latitude spots
- Can't be explained by solar dynamo

Roettenbacher et al. (2016)






# Spotted Stars



Roettenbacher et al. (2016)

# O-Star Triple Sigma Orionis



Schaefer et al. 2016

$M_{Aa} = 16.99 \pm 0.20 M_{\odot}$   
 $M_{Ab} = 12.81 \pm 0.18 M_{\odot}$   
 $d = 387.5 \pm 1.3 \text{ pc}$

- Dynamical masses for 3 O-stars
- Distance to sigma Orionis cluster
- Inner and outer orbits are not coplanar (120 – 127 deg)

# Interacting Binaries

Beta Lyrae


P = 13 days  
a = 0.87 mas



Zhao et al. (2008)

Algol


P = 2.9 days  
a = 2.2 mas



Baron et al. (2012)

# Transiting Disk: Epsilon Aurigae



## Epsilon Aurigae Eclipse (CHARA-MIRC)



Kloppenborg et al. (2010)







# Transiting Disk: Epsilon Aurigae



Kloppenborg et al. (2015)



# High Contrast Binaries



- Cepheid V1334 Cyg
- P = 5 yr
- Companion contributes 3.1% of flux
- Gallenne et al. (2013,2015)



# RS CVn Binaries




Binary orbit + ellipsoidal variations



Roettenbacher et al. (2015a, 2015b)

# Be Stars

- Rapidly rotating B-type stars that eject gas into a circumstellar disk
- Evidence for the disks
  - Rotationally broadened emission lines
  - IR excess
  - Linear polarization
  - Spatially resolved through interferometry
- Variable on time-scales of days to decades






# Be Stars



Geometry and physical structure of disks

Gies et al. (2007)


# Kinematic Model Be Stars



Mourard et al. 2015



# Binarity in Be Stars

- Role of binarity in Be stars – past mass transfer events?
  - Spun up secondary orbiting stripped down remnant companion (neutron star, white dwarf, helium star)
  - High contrast at close separations



Mourard et al. (2015)

# Disk wind in AB Aurigae




- Resolve H $\alpha$  formation region in young accreting intermediate mass star
- Bulk of H $\alpha$  forms in disk wind from innermost regions (0.05 – 0.15 AU)
- Perraut et al. (2016)





# Classical Nova

- Material from close binary companion accretes onto surface of white dwarf
- When pressure and temperature of accreted material reach a critical level, ignites in a thermonuclear runaway
- Expansion velocities of 500 – 3000 km/s




# Nova Delphini 2013



- Changes in apparent expansion – optically thick core surrounded by diffuse envelope that cools over time
- Geometric distance (4.5 kpc)
- Asymmetric shape detected as early as  $t = 2$  days

Schaefer et al. 2014





# Summary

- Exciting science opportunities
  - 146 refereed papers and counting
- AO + updated detectors + community input
- Many more years of productive science programs in the future