



CHARA Community Workshop:

Milliarcsecond Astronomy with the CHARA Array



Gail Schaefer

The CHARA Array of
Georgia State University

Mount Wilson, CA



CHARA Community Workshop


Schedule

Stellar Astrophysics at High Angular Resolution	Gail Schaefer	2:00 – 2:45 pm
Overview of the CHARA Array and Applying for Time	Douglas Gies	2:45 – 3:30 pm
Planning Observations, Data Access, and Software Tools	Gail Schaefer	3:30 – 4:00 pm
Open Discussion	All	4:00 – 5:00 pm

Thanks to David Soderblom and the staff at STScI for hosting us!



Angular Resolution





Long Baseline Optical/Infrared Interferometers



CHARA Array - Mount Wilson, CA



NPOI - Anderson Mesa, AZ



VLT Interferometer - Paranal, Chile




MROI - Magdalena Ridge, NM
(under construction)



Principles of Interferometry




Interferometer




Resolution ~ 0.5 mas for 300 meter baseline in the H-band ($1.6 \mu\text{m}$)

Fringe Visibility



- Amplitude of fringes = Visibility
 - Point Source: $V = 1.0$
 - Resolved source: loss of coherence reduces fringe visibility
 - Measures the size and geometry of source



Fringe Visibility



- The visibility is the Fourier Transform of the brightness distribution
- Analytic functions for simple geometries
- Berger & Segransan
“Introduction to visibility modeling” 2007, New Ast Rev, 51, 576






Angular Diameters



- Visibility amplitude
 - size and structure of source




Angular Diameters



- Visibility amplitude
 - size and structure of source




Surface Features



Roettenbacher et al. 2016



Surface Features




Roettenbacher et al. 2016



Binary Stars

Separated Fringe Packet Binaries




Farrington et al. (2010)




Binary Stars

Visibility Modulation



- Fringe packets for the two components overlap
- Fringe visibility varies periodically
 - binary separation
- Minimum in curve
 - flux ratio = $\frac{1 - V_{min}}{1 + V_{min}}$

Closure Phase




Monnier, "Phases in Interferometry" 2007, New Ast Rev, 51, 604

- Atmosphere corrupts phase information at vis/IR wavelengths
- Closure phase (3 or more telescopes):
 - $CP = \Phi_{12} + \Phi_{23} + \Phi_{31}$
- Cancels atmospheric effects
- Point symmetric object will have closure phase of 0° or 180°
- Measures asymmetries in source distribution




Binary Stars

Visibility (S1-E1)



Closure Phase (S1-E1-W1)





Flux ratio = 0.99





Binary Stars

Visibility (S1-E1)

Closure Phase (S1-E1-W1)




Flux ratio = 0.99



Flux ratio = 0.50

Differential Visibilities and Phases



- Spectrally dispersed interferometry
 - emission lines (BrG, Ha)
 - velocity structure

- Drop in visibility across emission line
 - variation in size and flux ratio between star and disk
- “S” shaped profile in differential phase
 - photo-center shift across wavelength channels




Interferometric Observables

- Visibility amplitude
 - size and structure of source
- Closure phase
 - asymmetries in source distribution
- Differential visibilities and phases
 - emission lines
 - velocity structure



UV Coverage

$$u = B_x / \lambda$$
$$v = B_y / \lambda$$





- Interferometer baseline projected on to plane of sky
- Position angle and projected baseline length will change as the earth rotates



UV Coverage

$$u = B_x / \lambda$$
$$v = B_y / \lambda$$




- Interferometer baseline projected on to plane of sky
- Position angle and projected baseline length will change as the earth rotates




Science at High Angular Resolution

Stellar Diameters




- Empirical HRD
- ~ 290 stars, $\sigma_{\theta} < 5\%$
- Angular diameter + parallax
 - Linear radius

Stellar Diameters





- Empirical HRD
- ~ 290 stars, $\sigma_{\theta} < 5\%$
- Angular diameter + parallax
 - Linear radius
- Effective Temperature

$$- F_{bol} = \frac{1}{4} \theta^2 \sigma T^4$$






Stellar Ages



- Comparison of R , T_{eff} , L with evolutionary models
 - Masses and ages of stars
- Age of the Ursa Major moving group: 414 ± 23 Myr
 - Diameters of A-stars (Jones et al. 2015)


Exoplanet Host Stars



- Age and mass of host star
- Size of habitable zones
 - L , T_{eff}
- Physical parameters of planets
 - Radius of transiting planets



Asteroseismology



Mass, radius, mean density, and surface gravity (need T_{eff})

$$v_{\text{max}} \propto (M / R^2) (T_{\text{eff}})^{-0.5}$$


$$\langle v_{\text{nl}} \rangle \propto \langle \rho \rangle^{0.5}$$

Oscillation power spectrum

$\langle v_{\text{nl}} \rangle$: frequency separation of modes

v_{max} : frequency of maximum power

Asteroseismology




Mass, radius, mean density, and surface gravity (need T_{eff})

$$\nu_{\text{max}} \propto (M / R^2) (T_{\text{eff}})^{-0.5}$$

$$\langle \nu_{\text{nl}} \rangle \propto \langle \rho \rangle^{0.5}$$

Oscillation power spectrum
 $\langle \nu_{\text{nl}} \rangle$: frequency separation of modes
 ν_{max} : frequency of maximum power



Test asteroseismic scaling relations for main sequence stars
 Huber et al. (2012)



Asteroseismology: Transiting Exoplanet Survey Satellite

- **TESS Input Catalog**
 - 596 million objects
 - 200,000 – 400,000 selected for high cadence
- **Two-year mission**
- **Launched on April 18, 2018**
- **$V < 7$ mag**
 - 4,864 stars resolvable ($\theta > 0.2$ mas)
- **$V < 8$ mag**
 - 13,922 stars resolvable ($\theta > 0.2$ mas)






Image credit: NASA


Cepheids



Merand et al 2005


- Radial velocity and angular diameter variation over pulsational phase
- Calibration of Baade-Wesselink technique - pulsation parallaxes
- Simultaneously fit photometry, spectroscopy, interferometry (Merand et al. 2015)
 - Mitigate systematics in projection factor
 - 2% accuracy on radius and distance

Rapid Rotators




- Oblateness
- Gravity darkening
 - $T_{\text{eff}} \sim g^{\beta}$

Rapid Rotators




- Oblateness
- Gravity darkening
 - $T_{\text{eff}} \sim g^{\beta}$
 - von Zeipel model: $\beta = 0.25$
 - empirically derived $\beta = 0.19$




Spotted Stars

2011



2013




Roettenbacher et al. (2016)

- Magnetically active star zeta Andromedae
- Rotation Period: 18 days
- $\theta = 2.502 \pm 0.008$ mas
- Direct confirmation of persistent polar spot
- Transient lower latitude spots
- Can't be explained by solar dynamo



Spotted Stars




Roettenbacher et al. (2016)




Binary Stars: Orbits and Dynamical Masses

Schaefer
et al. 2016



- Spatially resolved orbits of spectroscopic binaries
 - Masses and distances to 1-3% precision


Simon-Diaz
et al. 2015






Binary Stars: Orbits and Dynamical Masses

Schaefer
et al. 2016






Simon-Diaz
et al. 2015



- Spatially resolved orbits of spectroscopic binaries
 - Masses and distances to 1-3% precision
 - O-Star Triple sigma Orionis
 - $M_{Aa} = 16.99 \pm 0.20 M_{\odot}$
 - $M_{Ab} = 12.81 \pm 0.18 M_{\odot}$
 - $d = 387.5 \pm 1.3 \text{ pc}$
 - Distance to sigma Orionis cluster
- Schaefer et al. 2016

High Contrast Binaries




- High contrast companions
 - Separations 0.5 - 50 mas
 - $\Delta H < 6$ mag
 - Cepheids companions - Gallenne et al. 2013, 2015
 - RS CVn companions - Roettenbacher et al. 2015a, 2015b

Interacting Binaries

Beta Lyrae

$P = 13$ days
 $a = 0.87$ mas



- Mass donor is elongated – detection of photospheric distortion due to Roche lobe filling
- Thick disk around mass gainer - elongated


Zhao et al. (2008)



Interacting Binaries

Beta Lyrae


$P = 13$ days
 $a = 0.87$ mas



Zhao et al. (2008)

Algol


$P = 2.9$ days
 $a = 2.2$ mas



Baron et al. (2012)

Transiting Disk: Epsilon Aurigae



Epsilon Aurigae Eclipse (CHARA-MIRC)



Limb-Darkened Disk:
 $\theta_{LDD} = 2.22 \pm 0.09$ mas
 $\mu_{LDD} = 0.50 \pm 0.26$



Kloppenborg et al. (2010)

Transiting Disk: Epsilon Aurigae



Kloppenborg et al. (2015)



Be Stars



- Rapidly rotating B-type stars that eject gas into a circumstellar disk
- Geometry and physical structure of disks
- Kinematics
- Size vs. wavelength
- Investigate variability over time

Gies et al. (2007)


Kinematics of Be Star Disks



Mourard et al. 2015

Binarity in Be Stars


- Role of binarity in Be stars – past mass transfer events?
 - Spun up secondary orbiting stripped down remnant companion (neutron star, white dwarf, helium star)
 - High contrast at close separations






Mourard et al. (2015)

Classical Nova

- Material from close binary companion accretes onto surface of white dwarf
- When pressure and temperature of accreted material reach a critical level, ignites in a thermonuclear runaway
- Expansion velocities of 500 – 3000 km/s



Nova Delphini 2013




Schaefer et al. 2014

- Changes in apparent expansion – optically thick core surrounded by diffuse envelope that cools over time
- Geometric distance (4.5 kpc)
- Asymmetric shape detected as early as $t = 2$ days



Looking Toward the Future...



JMMC Stellar
Diameter Catalog


DEC > -20°
V < 8 mag
 $\theta > 0.2$ mas

Nstar = 18,147

Imaging = 9,781



Looking Toward the Future...



JMMC Stellar
Diameter Catalog

DEC > -20°
H < 8 mag
 $\theta > 0.5$ mas

Nstar = 19,116

Imaging = 3,558



Summary

- CHARA Array can resolve sizes of stars across the HR Diagram
- Improving our understanding of stellar structure and evolution
 - Stellar radius, effective temperature, dynamical masses
 - Limb darkening, gravity darkening
 - Rotation
 - Starspots
 - Mass loss
 - Convection
- Community access time
 - 50 nights available per year
 - NOAO proposals due in September and March